

Errata:

Better Lower Bounds on Detecting Affine and Spherical Degeneracies

Jeff Erickson
Computer Science Department
Duke University
Durham, NC 27708-0129
jeffe@cs.duke.edu
<http://www.cs.duke.edu/~jeffe>

Raimund Seidel
Fachbereich Informatik
Universität des Saarlandes
D-66123 Saarbrücken, Germany
seidel@cs.uni-sb.de

January 9, 1996

The lower bounds in [1] are based on the following adversary argument. Initially, the adversary presents a nondegenerate set of points. If an algorithm does not do enough work, the adversary can modify its original input, introducing a degeneracy that the algorithm cannot detect. To prove some of our lower bounds, we first construct an adversary input for a restricted problem, in which some degeneracies are ignored, and then argue that this point set can be perturbed into general position to obtain an adversary input for the unrestricted problem. This perturbation technique is based on an incorrect assumption, and the proofs that use it are invalid.

Let Φ be a collection of D -variate real polynomials. Each polynomial in Φ naturally induces an algebraic surface in \mathbb{R}^D , and these surfaces define a cellular decomposition of \mathbb{R}^D . Specifically, two points $p, q \in \mathbb{R}^D$ are in the same cell if and only if the sign of every polynomial in Φ is the same at both points. A surface σ *nicely bounds* a cell C in this decomposition if there is a continuous path from some point in C to some point outside C that intersects σ transversely and is either completely contained in or completely disjoint from every other surface induced by Φ .

The correctness of the perturbation technique is based on the following incorrect claim [1, p.48]:

Let C be an arbitrary cell, and let C' be one of the cells in its boundary. Then every surface that nicely bounds C' also nicely bounds C .

A counterexample is given by a set of four surfaces in \mathbb{R}^3 , defined by the polynomials $y - xz$, $x - yz$, $z - 1$ and $z + 1$. See Figure 1. Let C' be the line segment joining $(0, 0, -1)$ and $(0, 0, 1)$. C' is a cell in the decomposition defined by these polynomials, and it is nicely bounded by the planes $z = \pm 1$. However, no other cell is nicely bounded by both planes. In fact, none of the 2-dimensional cells that contain C' is nicely bounded by either plane.

Because of this error, the following results have incorrect proofs in [1]: Lemma 3.1, Theorem 3.2, Theorem 3.3, Theorem 4.2, Corollary 4.3, Theorem 5.2, Theorem 5.3, Theorem 5.6, and Corollary 5.7. Correct (and much simpler) proofs of all but one of these results appear in [3]. The only theorem whose proof we cannot correct is Theorem 5.6. Proving an $\Omega(n^{d+1})$ lower bound for arbitrary spherical degeneracies in \mathbb{R}^d remains an open problem.

A similar error appears in Section 5 of [2]. The proofs in that section are also invalid, although similar results are proven correctly in [3].




Figure 1: A counterexample to a claim in [1].

Acknowledgments We would like to thank one of the anonymous referees of the journal version of [2] for demanding a proof of the false claim.

References

- [1] J. Erickson and R. Seidel. Better lower bounds on detecting affine and spherical degeneracies. *Discrete Comput. Geom.* 13:41–57, 1995.
- [2] J. Erickson. Lower bounds for linear satisfiability problems. *Proc. 6th Ann. ACM-SIAM Symp. on Discrete Algorithms*, pp. 388–395, 1995.
- [3] J. Erickson. New lower bounds for convex hull problems in odd dimensions. *Proc. 12th Ann. ACM Symp. on Computational Geometry*, pp. 1–9, 1996.